

New approach on Supersymmetric Quantum Systems:Real and Complex Hamiltonians.

Biswanath Rath

Department of Physics, North Orissa University, Takatpur,
Baripada -757003, Odisha, INDIA
(e.mail:biswanathrath10@gmail.com)

We propose a new method for generating new Hamiltonians for the development of supersymmetric Quantum Mechanics for real as well as complex systems including **twins** and **quadruplets** Hamiltonians. Interestingly Hamiltonians, which can be realised in this method , can hardly be derived using all previous formalisms developed so far.Further examples discussed here include both shape invariant and shape non-invariant potentials .We notice a new thing (which has not been revealed earlier) in the case of shape invariant potentials i.e by suitable choice of parameter in superpotential

the generated Hamiltonians satisfy either (i) SUSY conditions i.e $E_n^{(+)} = E_{n+1}^{(-)}$; $E_0^{(-)} = 0$ or (ii) Iso-spectral condition i.e $E_n^{(+)} = E_n^{(-)}$. However for non-shape invariant potentials energy relations depend only on the choice of superpotentials. We present both analytical as well as numerical results.

PACS 11.30.Pb ,03.65Db,11.30.Er,03.65.Ge.

Key words-Superpotential,SUSY condition,Iso-spectral condition ,real Hamiltonians,complex Hamiltonians ,exactly solvable systems,numerical results.

I.Introduction

Generation of Quantum systems has been a subject of main interest since the early development of Quantum Mechanics. In fact it gained momentum after the work of Witten [1] ,who proposed that similar quantum systems can be generated through "Supersymmetry". However ,Gendeshstein [2] proposed that when generated Hamiltonians(H_+ , H_-) reflect similar

shape in potentials (commonly known as shape invariant) then energy eigenvalues can be related to each other obey only SUSY conditions i.e

$$E_n^{(+)} = E_{n+1}^{(-)} \quad (1)$$

and

$$E_0^{(-)} = 0 \quad (2)$$

where $E_n^{(+)} \rightarrow H_+$ and $E_n^{(-)} \rightarrow H_-$. Interestingly ,not only a report but also a book on supersymmetry has been published by Cooper,Khare and Sukhatme [3] where the authors strongly believe in the validity of relations in Eqs(1,2) and proposed many exactly solvable Hamiltonians reflecting shape invariants potentials i.e

$$V_+(x, a_0) = V_-(x, a_1) + R(a_1) \quad (3)$$

to justify the same.Mean while PT symmetric Quantum Mechanics has gained momentum particularly after the work of Bender and Boettecher [4] .In order to develop the supersymmetry in PT systems Bazeia,Das,Greenwood and

Lanso [5] proposed that for shape invariant potentials (as in Eq(3)) only Iso-spectral condition

$$E_n^{(+)} = E_n^{(-)} \quad (4)$$

is the most appropriate one .Of course authors [5] cite examples to justify their stand.Here we notice that when the generated Hamiltonians are Hermitian in nature they obey SUSY condition [3], however when generated Hamiltonians determined using siumilarity type of transformations [5] in PT symmetric systems ,obey Iso-spectral condition.Hence we feel there is lacking co-operation between two groups of authors [3,5] .Further from the literature on Supersymmetry we notice that Hamiltonian of the type

$$H = (p + ig)^2 + V(x) \quad (5)$$

proposed much earlier by Hatano and Nelson [6] ,has not been considered [5,3]. Hence from the above discussions we feel that existing work on Supersymmetry needs development.Here we propose a new method for the development of Supersymmetry in Quantum Systems which can be visualized in real or

complex Hamiltonians .Futher interesting examples(including exactly solvable shape invariant potentials) are presented in addition to numerical calculation.

II.Generators,Superpotentials and Hamiltonians.

Let us define two generators using operator theory as

$$A = -\frac{d}{dx} + iW(x) \quad (6)$$

$$B = \frac{d}{dx} + iW^*(-x) \quad (7)$$

where superpotential $W(x)$ can be real or complex.It may not

be practically possible to generate either A from B or

vice-versa.Now using A and B we define two Hamiltonians

$$H_+ = AB = p^2 - [W(x) - W^*(-x)]p - i\frac{dW^*(-x)}{dx} - W(x)W^*(-x) \quad (8)$$

and

$$H_- = BA = p^2 - [W(x) - W^*(-x)]p + i\frac{dW(x)}{dx} - W(x)W^*(-x) \quad (9)$$

One can notice that $AB \neq BA$ i.e $H_+ \neq H_-$. This type of generated Supersymmetric Hamiltonians are new to the literature and can not be visualised by any of the previous work[1,2,3,5].

IIA.SUSY conditions :Real Hamiltonian

Let us consider that

$$W(x) = ix \tag{10}$$

in this case we find

$$PTW(x)PT = W(x) = ix \tag{11}$$

The generated Hamiltonians are

$$H_- = p^2 + x^2 - 1 \tag{12}$$

and

$$H_+ = p^2 + x^2 + 1 \tag{13}$$

It is easy to see that the corresponding energy levels are

$$E_n^{(-)} = 2n \tag{14}$$

$$E_n^{(+)} = 2n + 2 = E_{n+1}^{(-)} \quad (15)$$

with

$$E_0^{(-)} = 0 \quad (16)$$

where $E_n^{(+)} \rightarrow H_+$ and $E_n^{(-)} \rightarrow H_-$.

IIB.SUSY conditions :Complex Hamiltonian

Now consider that

$$W(x) = ix - ig \quad (17)$$

In this case for $g \neq 0$ we find

$$PTW(x)PT \neq W(x) = ix - ig \quad (18)$$

The generated Hamiltonians are

$$H_- = (p + ig)^2 + x^2 - 1 \quad (19)$$

and

$$H_+ = (p + ig)^2 + x^2 + 1 \quad (20)$$

In order to calculate eigenvalues of the above Hamiltonians, we use $SU(11)$ algebra [6] . Let us consider the Hamiltonian

of the type [6]

$$H = h_{11}p^2 + h_{22}x^2 + ih_{12}(xp + px) + ih_1p + h_2x \quad (21)$$

having energy eigenvalue

$$\epsilon_n = \sqrt{h_{11}h_{22} + h_{12}^2}(2n + 1) + \frac{(h_1^2h_{22} - h_2^2h_{11} - 2h_1h_2h_{12})}{4(h_{11}h_{22} + h_{12}^2)} \quad (22)$$

Now using the above formula we find for H_- ; H_+ one has

(i) $h_1 = h_2 = 1$ (ii) $h_{12} = h_2 = 0$ (iii) $h_1 = 2g$. Hence it is easy to show that

$$E_n^{(-)} = 2n \quad (23)$$

and

$$E_n^{(+)} = 2n + 2 = E_{n+1}^{(-)} \quad (24)$$

with

$$E_0^{(-)} = 0 \quad (25)$$

where $E_n^{(+)} \rightarrow H_+$ and $E_n^{(-)} \rightarrow H_-$. This type of Hamiltonian was suggested earlier [7]. Interestingly we realise it in a

systematic way in the context of Supersymmetry and notice that SUSY is not broken . Interestingly the above Hamiltonian is non- $\mathcal{PT}$ invariant and non-Hermiticity in nature . Further to support the above example on SUSY condition , we consider another interesting form of $W(x)$ as

$$W(x) = ikx^3 - ix^2g \quad (26)$$

The corresponding Hamiltonians are the following

$$H_- = p^2 + 2igx^2p - 3kx^2 + 2gx - g^2x^4 + k^2x^6 \quad (27)$$

and

$$H_+ = p^2 + 2igx^2p + 3kx^2 + 2gx - g^2x^4 + k^2x^6 \quad (28)$$

Now using matrix diagonalisation method [8] we calculate the eigenvalues and reflect the same in table-1 **Table -1**

First five eigenvalues of SUSY Hamiltonians .

k	g	$E_n^{(-)}$	$E_n^{(+)}$
1	1	0	1.935 482
		1.935 482	6.298 495
		6.298 495	11.680 970
		11.680 970	18.042 635
		18.042 634	25.254 604
2	2	0	2.737 184
		2.737 184	8.907 417
		8.907 417	16.519 386
		16.519 389	25.516 139
		25.516 137	35.715 404

IIC.Iso-spectral condition :Complex Hamiltonian

In the previous examples we have considered superpotentials having complex in nature.Now we consider superpotentials having real in nature i.e

$$W(x) = x^2 \tag{29}$$

in this case we find

$$PTW(x)PT = W(x) = x^2 \quad (30)$$

The generated Hamiltonians are

$$H_- = p^2 - x^4 - 2ix \quad (31)$$

and

$$H_+ = p^2 - x^4 + 2ix \quad (32)$$

The energy eigenvalues of these Hamiltonians are obtained by solving Schroedinger's equation numerically [9] and are tabulated in Table-2. From the table one can see that energy levels satisfy the iso-spectral condition [7,8] i.e

$$E_n^{(+)} = E_n^{(-)} \quad (33)$$

Table -2

First five eigenvalues of Iso-spectral Hamiltonians .

Level	$E_n^{(+)}$	$E_n^{(-)}$
0	0	0
1	3.398 150	3.398 150
2	8.700 453	8.700 453
3	14.977 808	14.977 808
4	21.999 001	21.999 001

**IID.SUSY conditions: Exactly solable shape
invariant Potentials.**

**Let us consider one interesting type of complex
superpotential as**

$$W(x) = ix - i\frac{\lambda}{x} \quad (34)$$

in this case we find

$$PTW(x)PT = W(x) = ix - i\frac{\lambda}{x} \quad (35)$$

**The present superpotential contains a linear term
and an inverse term ,whose similar type was first
reflected in the work of Supersymmetry by
Rath[9] in the context of Hermitian**

Hamiltonian. Later on this type was used by Bazeia et.al [5] by making a modification as

$$W(x) = ix + \epsilon - \frac{a_0}{(ix + \epsilon)} \quad (36)$$

in the context of PT symmetry of supersymmetric Quantum Mechanics to discuss iso-spectral conditions. Interestingly Bazeia et.al [5] have not reflected neither numerical nor analytical results to justify their stand. Here the generated Hamiltonians are obtained using slight modified form of superpotential as

$$H_+ = \frac{p^2}{2} + \frac{x^2}{2} + \frac{\lambda(\lambda + 1)}{2x^2} - \lambda + 0.5 \quad (37)$$

and

$$H_- = \frac{p^2}{2} + \frac{x^2}{2} + \frac{\lambda(\lambda - 1)}{2x^2} - \lambda - 0.5 \quad (38)$$

It is easy to see that the corresponding exact energy levels are

$$E_n^{(-)} = 2n + 0.5 + 0.5\sqrt{1 + 4\lambda(\lambda - 1)} - \lambda \quad (39)$$

$$E_n^{(+)} = 2n + 1.5 + 0.5\sqrt{1 + 4\lambda(\lambda + 1)} - \lambda \quad (40)$$

One can see that for any value of $\lambda > 2$,the energy levels satisfy the SUSY conditions as

$$E_n^{(+)}(\lambda > 2) = 2n + 2 \quad (41)$$

$$E_n^{(-)}(\lambda > 2) = 2n \quad (42)$$

with

$$E_0^{(-)}(\lambda > 2) = 0 \quad (43)$$

Interesting point is that eigenvalues are independent of λ . The corresponding wave functions are the following:

$$\Psi_n^{(+)} = C_n^+ \left[\left(-\frac{d}{dx} + x \right)^2 - \frac{\lambda(\lambda + 1)}{x^2} \right]^n x^{0.5 + \sqrt{0.25 + \lambda(\lambda + 1)}} e^{-x^2/2} \quad (44)$$

and

$$\Psi_n^{(-)} = C_n^- \left[\left(-\frac{d}{dx} + x \right)^2 - \frac{\lambda(\lambda - 1)}{x^2} \right]^n x^{0.5 + \sqrt{0.25 + \lambda(\lambda - 1)}} e^{-x^2/2} \quad (45)$$

II E. Iso-spectral condition: Exactly solable shape invariant Potentials.

Here we consider slight modification of above
complex superpotential as

$$W(x) = ix + i\frac{\lambda}{x} \quad (46)$$

in this case we find

$$PTW(x)PT = W(x) = ix + i\frac{\lambda}{x} \quad (47)$$

Here the generated Hamiltonians are obtained
using slight modified form of superpotential as

$$H_+ = \frac{p^2}{2} + \frac{x^2}{2} + \frac{\lambda(\lambda - 1)}{2x^2} + \lambda + 0.5 \quad (48)$$

and

$$H_- = \frac{p^2}{2} + \frac{x^2}{2} + \frac{\lambda(\lambda + 1)}{2x^2} + \lambda - 0.5 \quad (49)$$

It is easy to see that the corresponding exact
energy levels are

$$E_n^{(-)} = 2n + 0.5 + 0.5\sqrt{1 + 4\lambda(\lambda + 1)} + \lambda \quad (50)$$

$$E_n^{(+)} = 2n + 1.5 + 0.5\sqrt{1 + 4\lambda(\lambda - 1)} + \lambda \quad (51)$$

One can see that for any value of $\lambda > 2$,the energy levels satisfy the iso-spectral condition as

$$E_n^{(+)}(\lambda > 2) = E_n^{(-)}(\lambda > 2) = 2n + 2\lambda + 1 \quad (52)$$

Here energy levels are parameter dependent.

The corresponding wave functions are the following:

$$\Phi_n^{(+)} = D_n^+ \left[\left(-\frac{d}{dx} + x \right)^2 - \frac{\lambda(\lambda - 1)}{x^2} \right]^n x^{0.5 + \sqrt{0.25 + \lambda(\lambda - 1)}} e^{-x^2/2} \quad (53)$$

and

$$\Phi_n^{(-)} = D_n^- \left[\left(-\frac{d}{dx} + x \right)^2 - \frac{\lambda(\lambda + 1)}{x^2} \right]^n x^{0.5 + \sqrt{0.25 + \lambda(\lambda + 1)}} e^{-x^2/2} \quad (54)$$

III. Generators, Superpotentials : Twins and Quadruplets Hamiltonians.

In the above , we have generated only a pair of SUSY or Iso-spectra using suitable choice of superpotential . However in this generation we try to generate twin SUSY and Quadruplet iso-spectra

using the method given below . Let us define two generators using operator theory as

$$A = -ip - W_1 \quad (55)$$

$$B = ip - W_2 \quad (56)$$

such that the product W_1W_2 is an even function of x .

$$H_+ = AB = p^2 + i[W_2 - W_1]p + \frac{dW_2}{dx} + W_1W_2 \quad (57)$$

and

$$H_- = BA = p^2 + i[W_2 - W_1]p - \frac{dW_1}{dx} + W_1W_2 \quad (58)$$

One can notice that $AB \neq BA$ i.e $H_+ \neq H_-$. This type of generated Supersymmetric Hamiltonians are new to the literature and can not be visualised by any of the previous work[1,2,3,5]. Here as per the condition stated above one has to select $W_1; W_2$ even or odd .

vspace0.2in

IIIA. Twins SUSY

Case-(a) :: $W_1 = x$ and $W_2 = x^3$

$$H_1^+ = AB = p^2 + ix^3p - ixp + 3x^2 + x^4 \quad (59)$$

$$H_1^- = BA = p^2 + ix^3p - ixp - 1 + x^4 \quad (60)$$

Case-(b) :: $W_2 = x$ and $W_1 = x^3$

$$H_2^+ = AB = p^2 - ix^3p + ixp + 1 + x^4 \quad (61)$$

$$H_1^+ = AB = p^2 - ix^3p + ixp - 3x^2 + x^4 \quad (62)$$

In order to show the twin behaviour , we calculate the energy eigenvalues and reflect the same in table-3 using the matrix diagonalisation method [8].

Table - 3 :First four eigenvalues of Twin-SUSY Hamiltonians .

H_1^+	H_1^-	H_2^+	H_2^-
2.067 999 2	0	2.067 999 2	0
5.631 827 3	2.067 999 2	5.631 827 3	2.067 999 2
9.995 229 9	5.631 827 3	9.995 229 9	5.631 827 3
15.047 560 1	9.995 229 9	15.047 560 1	9.995 229 9

From the table one can see that H_1^+ and H_2^+ have the same eigenspectra even though the corresponding Hamiltonians are not the same . Similarly H_1^- and H_2^- pertain to same eigen spectra .Hence they are called twin-SUSY. It is worth to mention that the generated Hamiltonians are complex and satisfy $\mathcal{PT}$ - symmetry condition.

IIIB- Quadruplet Iso-Spectra

Case-(c) :: $W_1 = x^2$ and $W_2 = x^4$

$$H_3^+ = AB = p^2 + ix^4p - ix^2p + 4x^3 + x^6 \quad (63)$$

and

$$H_3^- = BA = p^2 + ix^4p - ix^2p - 2x + x^6 \quad (64)$$

Case-(d) :: $W_2 = x^2$ and $W_1 = x^4$

$$H_4^+ = AB = p^2 - ix^4p + ix^2p + 2x + x^6 \quad (65)$$

and

$$H_4^+ = AB = p^2 - ix^4p + ix^2p - 4x^3 + x^6 \quad (66)$$

In order to show the quadruplet behaviour , we calculate the energy eigenvalues and reflect the same in table-4 using the matrix diagonalisation method [8].

Table - 4 :First four eigenvalues of Quadruplet Iso-spectral Hamiltonians .

H_3^+	H_3^-	H_4^+	H_4^-
0.537 037 9	0.537 037 9	0.537 037 9	0.537 037 9
4.006 022 7	4.006 022 7	4.007 022 7	4.006 022 7
9.019 924 8	9.019 924 8	9.019 924 8	9.019 924 8
15.215 167 0	15.215 167 0	15.215 167 0	15.215 167 0

In this case the generated Hamiltonians are
non- $\mathcal{PT}$ invariant in nature .

IV.Conclusion.

In this paper ,we have developed a new method for supersymmetric Quantum System to realise it in real or complex operators.For shape invariant potentials we show that energy levels are intimately related reflecting susy or iso-spectral behaviour only on selection of parameters.Hence we co-relate previously published two different concepts [3,5,10] . We also generate two-Twins satisfying SUSY behaviour and quadruplet satisfying iso-spectral behaviour .It should be

remembered that present selections of A and B are not the only possibility . In fact one can follow this procedure to generate other types . In order to help the reader on new generation , we present another type as follows:

$$A_1 = i \frac{d}{dx} + W(x) \quad (67)$$

$$B_1 = i \frac{d}{dx} + W^*(-x) \quad (68)$$

where superpotential $W(x)$ can be real or complex. It may not be practically possible to generate either A_1 from B_1 or vice-versa. Now using

A_1 and B_1 we define two Hamiltonians

$$H_+ = A_1 B_1 = p^2 - [W(x) + W^*(-x)]p + i \frac{dW^*(-x)}{dx} + W(x)W^*(-x) \quad (69)$$

and

$$H_- = B_1 A_1 = p^2 - [W(x) + W^*(-x)]p + i \frac{dW(x)}{dx} + W(x)W^*(-x) \quad (70)$$

This new form can easily generate new type of SUSY Hamiltonians discussed recently by Marqques,Neg1eni and da Silva [12] . Here we select the form of $W(x)$ as

$$W(x) = i|x|^2 \quad (71)$$

and generate SUSY Hamiltonians as

$$H_- = p^2 - 2|x| + x^4 \quad (72)$$

$$H_+ = p^2 + 2|x| + x^4 \quad (73)$$

Further using matrix diagonalisation method , we present first five eigenvalues in table-5.

Table - 5

First four eigenvalues of New SUSY Hamiltonians and previous results [12] .

$E_n^{(+)}$	$E_n^{(-)}$	$E_n^{(+)}$ Previous [12]	$E_n^{(-)}$ Previous [12]
1.969 9	0	1.972 3	0
5.507 1	1.969 5	5.510 0	1.969 6
9.394 5	5.506 8	9.415 2	5.508 4
13.858 3	9.394 2	13.893 6	9.398 6

Hence we believe that present development will enrich the existing knowledge on supersymmetry in the context of energy calculation.

References

- [1] E.Witten Nucl.Phys ,B185,513(1981)
- [2] L.Gendenshtein Pis'ma Zh Eksp Teor Fiz 38 ,299 (1983).
- [3] F.Cooper,A.Khare and U.Sukhatme "Supersymmetry in Quantum Mechanics" (World Scientific,Singapore) (2001);Physics Reports 251(5,6)(1995).
- [4] C.M. Bender and S.Boettcher ,.Phys.Rev.Lett 80 , 5253 (1998).
- [5] D.Bazeia ,A.Das,L.Greenwood and L.Losano Phys. Lett. B673, 283 (2009).

- [6] H.B.Zhang
, G.Y.Jiang and G.C.Wang , J.Math.Phys. 56,
072103(2015)
- [7] N.Hatano and D.R.Nelson ,Phys.Rev.Lett. 77,
570(1996).
- [8] B.Rath,P.Mallick and P.K.Samal African
.Rev.Phys 9(0027, 201 (2014).
- [9] B.Rath,A.Nanayakkara,P.Mallick and
P.K.Samal (unpublished).
- [10]
B.Rath,ILNuovoCimentoB115,1229(2000);B124
N.8,1(2009).
- [11] A.M.Perelomov,Generalised Coherent
States and Their Applications (Springer Ver-
lag,Berlin)(1986).
- [12] F.Marques,O.Negrini and A.J.da Silva
,J.Phys.A45,115307(2012).